

A note on traveling wave solutions to the two component Camassa-Holm equation

K.Mohajer

Department of Mathematics and Statistics
University of Saskatchewan
106 Wiggins Road
Saskatoon, SK, S7N 5E6 CANADA
mohajer@math.usask.ca

March 12, 2008

Abstract

In this paper we show that non-smooth functions which are distributional traveling wave solutions to the two component Camassa-Holm equation are distributional traveling wave solutions to the Camassa-Holm equation provided that the set $u^{-1}(c)$, where c is the speed of the wave, is of measure zero. In particular there are no new peakon or cuspon solutions beyond those already satisfying the Camassa-Holm equation. However, the two component Camassa-Holm equation has distinct from Camassa-Holm equation smooth traveling wave solutions as well as new distributional solutions when the measure of $u^{-1}(c)$ is not zero. We provide examples of such solutions.

Mathematics Subject Classification. 35Q35, 35Q53

Keywords. Camassa-Holm equation, Traveling Waves, Peakons

The Camassa-Holm equation [1]

$$u_t + \kappa u_x - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx}, \quad (0.1)$$

arises as a model for the unidirectional propagation of shallow water waves over a flat bottom, $u(x, t)$ representing the water's free surface, and $\kappa \in \mathbb{R}$ being a parameter related to the critical shallow water speed. Camassa and Holm [1] discovered that the equation has non-smooth solitary waves that retain their individual characteristics through the interaction and eventually emerge with their original shapes and speeds. The traveling wave solutions of the Camassa-Holm equation have been classified by J. Lenells [4]. An alternative, and useful for generalizations form of this equation is

$$m_t + um_x + 2mu_x = 0, \quad (0.2)$$

where $m = u - u_{xx} + \frac{1}{2}\kappa$.

One such a generalization has been introduced by M. Chen, S. Liu and Y. Zhang [2]:

$$\begin{cases} m_t + um_x + 2mu_x - \rho\rho_x = 0, \\ \rho_t + (\rho u)_x = 0, \end{cases} \quad (0.3)$$

The traveling wave solutions are obtained by setting $u = u(x - ct)$ and $\rho = \rho(x - ct)$. In this case, easy manipulations show that (0.3) can be written as follows

$$\begin{cases} -2c(u' - u''') + 2\kappa u' + 3(u^2)' + ((u')^2)' - (u^2)''' = (\rho^2)', \\ -c\rho' + (\rho u)' = 0. \end{cases} \quad (0.4)$$

These equations are valid in the sense of distributions, if $u \in H_{loc}^1(\mathbb{R})$ and $\rho \in L_{loc}^2(\mathbb{R})$. Indeed, for a given function ρ , if $(\rho^2)' \in \mathcal{D}'(\mathbb{R})$, then $\rho \in L_{loc}^2(\mathbb{R})$. Since every distribution has a primitive which is a distribution (see [3]), we can integrate and then rewrite

$$\begin{cases} (v^2)'' = (v')^2 + p(v) - \rho^2, \\ \rho v = B_1. \end{cases} \quad (0.5)$$

where $v = u - c$ and $p(v) = 3v^2 + (2\kappa + 4c)v + K$ for some constants K and B_1 .

Definition 0.1. A pair of functions (u, ρ) where $u \in H_{loc}^1(\mathbb{R})$ and $\rho \in L_{loc}^2(\mathbb{R})$, is called a traveling wave solution for (0.3) if u and ρ satisfy (0.5) in the sense of distributions.

The following Lemma is due to J. Lenells [4].

Lemma 0.1. Let $p(v)$ be a polynomial with real coefficient. Assume that $v \in H_{loc}^1(\mathbb{R})$ satisfies

$$(v^2)'' = (v')^2 + p(v) \quad \text{in } \mathcal{D}'(\mathbb{R}). \quad (0.6)$$

Then

$$v^k \in C^j(\mathbb{R}) \quad \text{for } k \geq 2^j. \quad (0.7)$$

In our case, we have the following generalization:

Lemma 0.2. Let $p(v)$ be a polynomial with real coefficients. Assume that $v \in H_{loc}^1(\mathbb{R})$ and $\rho \in L_{loc}^2(\mathbb{R})$ satisfy the following system in $\mathcal{D}'(\mathbb{R})$:

$$\begin{cases} (v^2)'' = (v')^2 + p(v) - \rho^2, \\ \rho v = B_1. \end{cases} \quad (0.8)$$

Then

$$v^k \in C^j(\mathbb{R}) \quad \text{for } k \geq 2^j \quad \text{and } j \geq 0. \quad (0.9)$$

Proof. Since $v \in H_{loc}^1(\mathbb{R})$ and $\rho \in L_{loc}^2(\mathbb{R})$, (0.8) implies that $(v^2)'' \in L_{loc}^1(\mathbb{R})$. Therefore, $(v^2)'$ is absolutely continuous and $v^2 \in C^1(\mathbb{R})$. Also, since $v \in H_{loc}^1(\mathbb{R})$, then v is absolutely continuous and we can claim

$$(v^k)' = \frac{k}{2}(v^{k-2}(v^2)') \quad \text{for } k \geq 3.$$

To see why the claim is true, we first note that in fact, it is obviously true if k is an even number. Also, note that since the first derivative of an absolutely continuous function exists almost everywhere, in taking the first derivative of the product of two absolutely continuous functions we can use the Leibniz Rule almost everywhere. Now, if k is an odd number, let's say $k = 2n + 1$, then we can write

$$\begin{aligned} (v^k)' &= (v^{2n}v)' = v(v^{2n})' + v'v^{2n} \\ &= v(nv^{2(n-1)})(v^2)' + \frac{1}{2}(v^2)'v^{2n-1} \\ &= \frac{k}{2}v^{k-2}(v^2)'. \end{aligned}$$

Thus, we have

$$\begin{aligned} (v^k)'' &= \frac{k}{2}(v^{k-2}(v^2)')' \\ &= \frac{k}{2}((v^{k-2})'(v^2)' + v^{k-2}(v^2)'') \\ &= k(k-2)v^{k-2}(v')^2 + \frac{k}{2}v^{k-2}(v^2)'' \quad \text{for } k \geq 3. \end{aligned}$$

Substituting from (0.8) we have

$$\begin{aligned} (v^k)'' &= k(k-2)v^{k-2}(v')^2 + \frac{k}{2}v^{k-2}((v')^2 + p(v) - \rho^2) \\ &= k(k - \frac{3}{2})v^{k-2}(v')^2 + \frac{k}{2}v^{k-2}p(v) - \frac{k}{2}B_1v^{k-3}\rho. \end{aligned} \tag{0.10}$$

For $k \geq 3$ the right hand side of the above equation belongs to $L_{loc}^1(\mathbb{R})$. Therefore

$$v^k \in C^1(\mathbb{R}) \quad \text{for } k \geq 2. \tag{0.11}$$

Thus, the assertion holds for $j = 1$. We proceed by induction on j . Suppose

$$v^k \in C^{j-1}(\mathbb{R}) \quad \text{for } k \geq 2^{j-1} \quad \text{and } j \geq 2.$$

Then for $k \geq 2^j$ we have

$$\begin{aligned} v^{k-2}(v')^2 &= \frac{1}{2^{j-1}}(2^{j-1}v^{2^{j-1}-1}v') \frac{1}{k-2^{j-1}}((k-2^{j-1})v^{k-2^{j-1}-1}v') \\ &= \frac{1}{2^{j-1}(k-2^{j-1})}(v^{2^{j-1}})'(v^{k-2^{j-1}})' \in C^{j-2}(\mathbb{R}). \end{aligned} \tag{0.12}$$

Also, we have $v^{k-2}p(v) \in C^{j-1}(\mathbb{R})$ and $v^{k-3}\rho = B_1v^{k-4} \in C^{j-2}(\mathbb{R})$. Therefore the right hand side of equation (0.10) belongs to $C^{j-2}(\mathbb{R})$. Hence,

$$v^k \in C^j(\mathbb{R}) \quad \text{for } k \geq 2^j.$$

□

Remark. Lemma (0.2) implies that v' is possibly discontinuous only at points where $v = 0$. In fact, a much stronger result is true:

Corollary 0.1. *If $v \in H_{loc}^1(\mathbb{R})$ and $\rho \in L_{loc}^2(\mathbb{R})$ satisfy (0.8) in $\mathcal{D}'(\mathbb{R})$, then*

$$v \in C^\infty(\mathbb{R} \setminus v^{-1}(0))$$

and

$$\rho \in C^\infty(\mathbb{R} \setminus v^{-1}(0)).$$

Proof. Suppose $k \geq 2$. Then $v^k \in C^1(\mathbb{R})$. Therefore

$$kv^{k-1}v' = (v^k)' \in C(\mathbb{R}).$$

This implies that $v' \in C(\mathbb{R} \setminus v^{-1}(0))$. Thus, $v \in C^1(\mathbb{R} \setminus v^{-1}(0))$.

Now, assume that $v \in C^j(\mathbb{R} \setminus v^{-1}(0))$ for $j \geq 1$. For $k \geq 2^{j+1}$, we have $v^k \in C^{j+1}(\mathbb{R})$. Therefore

$$kv^{k-1}v' = (v^k)' \in C^j(\mathbb{R}).$$

This shows that $v' \in C^j(\mathbb{R} \setminus v^{-1}(0))$. Hence, $v \in C^{j+1}(\mathbb{R} \setminus v^{-1}(0))$. Thus, u is in the desired space. Now the statement for ρ follows from the second equation of (0.5). □

Remark. Since $v = u - c$, Corollary (0.1) shows that $u \in C^\infty(\mathbb{R} \setminus u^{-1}(c))$.

Since $\mathbb{R} \setminus u^{-1}(c)$ is an open set, we have

$$\mathbb{R} \setminus u^{-1}(c) = \bigcup_{i=1}^{\infty} (a_i, b_i).$$

So, u is smooth in every interval (a_i, b_i) where the following Lemma holds (below $(a_i, b_i) = (a, b)$):

Lemma 0.3. *Let (u, ρ) be a traveling wave solution to (0.3). Suppose u is smooth in the interval (a, b) . Then in the interval (a, b) , u satisfies the following equation:*

$$(u - c)^2 u'^2 = P(u), \tag{0.13}$$

where

$$P(u) = (u^2 + \kappa u + A)(u - c)^2 + C(u - c) + B, \tag{0.14}$$

and A, B and C are some constants.

Proof. Since both u and ρ are smooth in (a, b) we use standard calculus rules. By the first equation of (0.5), we have

$$2(v')^2 + 2vv'' = (v')^2 + p(v) - \rho^2.$$

Therefore,

$$(v')^2 + 2vv'' = p(v) - \rho^2.$$

Multiplying by v' we have

$$(v')^3 + v((v')^2)' = v'p(v) - v'\rho^2.$$

Thus,

$$(v(v')^2)' = v'p(v) - v'\rho^2.$$

Hence,

$$(v(v')^2)' = (3v^2 + (2\kappa + 4c)v + K)v' - \frac{Bv'}{v^2},$$

where $B = B_1^2$.

Integration yields

$$v(v')^2 = v^3 + (\kappa + 2c)v^2 + Kv + \frac{B}{v} + C.$$

Now, multiplying this equation by v we get

$$v^2(v')^2 = (v^2 + (\kappa + 2c)v + K)v^2 + Cv + B.$$

Substituting $v = u - c$ and simplifying, we have

$$(u - c)^2(u')^2 = (u^2 + \kappa u + A)(u - c)^2 + C(u - c) + B,$$

for some constant A . □

Theorem 0.1. *Suppose (u, ρ) is a non-smooth traveling wave solution to (0.3). If $u^{-1}(c)$ is a set of measure zero, then u is a solution to the Camassa-Holm equation.*

Proof. Suppose $\xi \in \mathbb{R} \setminus u^{-1}(c)$. Since, $u^{-1}(c) \neq \emptyset$, there exists an $\eta \in u^{-1}(c)$ such that either $\xi > \eta$ or $\xi < \eta$. Without loss of generality, assume that $\xi < \eta$. Let $\eta_0 = \inf\{\eta \in u^{-1}(c) : \eta > \xi\}$. Since $u^{-1}(c)$ is a closed set, $\eta_0 \in u^{-1}(c)$. So, $(\xi, \eta_0) \subseteq \mathbb{R} \setminus u^{-1}(c)$. Thus, we have proved that there exists an $\eta \in u^{-1}(c)$ such that either $(\xi, \eta) \subseteq \mathbb{R} \setminus u^{-1}(c)$ or $(\eta, \xi) \subseteq \mathbb{R} \setminus u^{-1}(c)$. Now, consider the equation (0.13) and set $F(u) = \frac{P(u)}{(u-c)^2}$. We claim that B in (0.13) equals 0. Suppose $B \neq 0$. Since $B = B_1^2$, we have $B > 0$. Then (0.14) implies that

$$\frac{1}{\sqrt{F(u)}} = \frac{1}{\sqrt{B}}|u - c| + \mathcal{O}((u - c)^2) \quad u \rightarrow c.$$

On the other hand, we have

$$\frac{d\xi}{du} = \pm \frac{1}{\sqrt{F(u)}}.$$

Since $u \in C(\mathbb{R})$, for ξ close enough to η , integration yields

$$|\xi - \eta| = \frac{1}{2\sqrt{B}}(u - c)^2 + \mathcal{O}((u - c)^3) \quad u \rightarrow c. \quad (0.15)$$

Therefore,

$$|\xi - \eta| = \frac{1}{2\sqrt{B}}(u - c)^2(1 + \mathcal{O}(u - c)) \quad u \rightarrow c.$$

So,

$$|\xi - \eta|^{\frac{1}{2}} = \frac{1}{\sqrt{2\sqrt{B}}} |u - c| \sqrt{1 + \mathcal{O}(u - c)} \quad u \rightarrow c.$$

Thus,

$$|\xi - \eta|^{\frac{1}{2}} = \frac{1}{\sqrt{2\sqrt{B}}} |u - c| (1 + \mathcal{O}(u - c)) \quad u \rightarrow c.$$

Hence,

$$|\xi - \eta|^{\frac{1}{2}} = \frac{1}{\sqrt{2\sqrt{B}}} |u - c| + \mathcal{O}((u - c)^2) \quad u \rightarrow c.$$

This implies that

$$(u - c) = \mathcal{O}((\xi - \eta)^{\frac{1}{2}}) \quad \xi \rightarrow \eta.$$

Therefore,

$$(u - c)^2 = \mathcal{O}(\xi - \eta) \quad \xi \rightarrow \eta.$$

Thus, we have

$$|u - c| = \sqrt{2\sqrt{B}} |\xi - \eta|^{\frac{1}{2}} + \mathcal{O}(\xi - \eta) \quad \xi \rightarrow \eta. \quad (0.16)$$

Hence,

$$\begin{aligned} |\xi - \eta|^{-\frac{1}{2}} - \sqrt{2\sqrt{B}} |u - c|^{-1} &= \mathcal{O}\left(\frac{|\xi - \eta|^{\frac{1}{2}}}{u - c}\right) \\ &= \mathcal{O}(1) \quad \xi \rightarrow \eta. \end{aligned}$$

So,

$$|u - c|^{-1} = \frac{1}{\sqrt{2\sqrt{B}}} |\xi - \eta|^{-\frac{1}{2}} + \mathcal{O}(1) \quad \xi \rightarrow \eta. \quad (0.17)$$

On the other hand, from (0.14) we have

$$|u'| = \sqrt{B}(u - c)^{-1} + \mathcal{O}(1) \quad \xi \rightarrow \eta. \quad (0.18)$$

Now combining (0.18) and (0.17), we have

$$|u'| = \frac{\sqrt[4]{B}}{\sqrt{2}} |\xi - \eta|^{-\frac{1}{2}} + \mathcal{O}(1) \quad \xi \rightarrow \eta. \quad (0.19)$$

Hence, $u' \notin L^2_{loc}(\mathbb{R})$. This contradiction shows that $B = 0$. Therefore, the second equation of (0.5) implies that $\rho = 0$ almost everywhere. $\square$

Now, we provide an example of a smooth solution of (0.3) that is not a solution of Camassa-Holm equation.

Example. Let $P(u)$ be as in the previous Theorem. Observe that $P(u) = (u - G)^2(u - L)^2$ if and only if

$$\begin{cases} \kappa = 2(c - (L + G)), \\ A = 2c\kappa - c^2 + (L + G)^2 + 2LG, \\ C = 2cA - \kappa c^2 - 2LG(L + G), \\ B = Cc - Ac^2 + L^2G^2. \end{cases} \quad (0.20)$$

Suppose $|u| < 1$ and $c > 1$. Therefore, if $G = -1$ and $L = 1$, integration yields

$$(1 - u)^{1-c}(1 + u)^{1+c} = e^{2(\xi - \xi_0)}. \quad (0.21)$$

Let's say $c = 2$ and $\xi_0 = 0$. We observe that the equation

$$\frac{(1 + u)^3}{1 - u} = e^{2\xi},$$

provides a smooth solution of (0.3) which is not a solution of Camassa-Holm equation. See figure 1.




Figure 1: (u on the left and ρ on the right) A smooth solution of (0.3) which is not a solution of Camassa-Holm equation.

The following Lemma provides necessary and sufficient conditions for a piecewise smooth function to be a distributional solution to (0.3).

Lemma 0.4. *Suppose u is a piecewise smooth function. The pair (u, ρ) is a distributional solution to (0.3) in the sense of definition 0.1 if and only if all of the following conditions hold:*

1. $u \in H_{loc}^1(\mathbb{R})$ and $\rho \in L_{loc}^2(\mathbb{R})$.
2. $(u - c)^2 \in W_{loc}^{2,1}(\mathbb{R})$.
3. u and ρ satisfy the equation (0.5) locally with the same constant K on every interval where u is smooth.

Proof. The part $(\Rightarrow)$ is easy. For the converse $(\Leftarrow)$, we note that since $(u - c)^2 \in W_{loc}^{2,1}(\mathbb{R})$, then $((u - c)^2)'$ is absolutely continuous and has no jumps. Therefore, $((u - c)^2)''$ defines a regular distribution [3]. Thus, every term in the equation (0.5) can be represented by an integral that defines a distribution on the space of test functions and we are allowed to write each integral as a finite sum of integrals over local intervals and use condition 3 to prove that u and ρ satisfy (0.5) in the sense of distributions. $\square$

Remark. We note that if the measure of $u^{-1}(c)$ is not zero, then the equation (0.5) implies that $\rho^2 = K$ on $u^{-1}(c)$. However in the Camassa-Holm equation if the measure of $u^{-1}(c)$ is not zero, then $K = 0$ because $\rho = 0$. This implies that solutions of the form given in the following example cannot arise from the Camassa-Holm equation.

Example. Set $\kappa = 0$. The pair of functions (u, ρ) given by

$$u(x) = \begin{cases} ce^{1-|x|} & \text{if } |x| > 1, \\ c & \text{if } |x| < 1, \end{cases}$$

$$\rho(x) = \begin{cases} c & \text{if } |x| < 1, \\ 0 & \text{if } |x| > 1, \end{cases}$$

is a solution to (0.3) but u is not a solution of Camassa-Holm equation. To see this, observe that the left hand side derivative of u at -1 and the right hand side derivative of u at 1 are non-zero and finite in contrast with the Camassa-Holm equation for which Lenells [4] showed that if the measure of $u^{-1}(c)$ is not zero, then these limits cannot be finite. See figure 2.




Figure 2: $u(x)$ is a solution of (0.3) but it is not a solution of Camassa-Holm equation.

Definition 0.2. Suppose f is a continuous function on $\mathbb{R}$.

1. We say f has a peak at x if f is smooth locally on both sides of x and

$$0 \neq \lim_{y \downarrow x} f'(y) = -\lim_{y \uparrow x} f'(y) \neq \pm\infty.$$

Traveling wave solutions of (0.3) with peaks are called peakons.

2. We say f has a cusp at x if f is smooth locally on both sides of x and

$$\lim_{y \downarrow x} f'(y) = -\lim_{y \uparrow x} f'(y) = \pm\infty.$$

Traveling wave solutions of (0.3) with cusps are called cuspons.

3. We say that f has a stump if there is an interval $[a, b]$ on which f is a constant and f is smooth locally to the left of a and to the right of b and

$$0 \neq \lim_{x \uparrow a} f'(x) = -\lim_{x \downarrow b} f'(x).$$

Traveling wave solutions of (0.3) with stumps are called stumpons. Note that, in the definition of a stump the limits can be either finite or infinite.

Theorem 0.1 limits the existence of new distributional peakon or cuspon solutions to the (0.3).

Corollary 0.2. Every peakon or cuspon traveling wave solution to (0.3) is a traveling wave solution to the Camassa-Holm equation.

Finally we would like to comment on the peaked solution reported in [2]. For reasons explained below, that solution is not a distributional solution. First, we note that by Corollary (0.1) the non-smooth points of a distributional solution

u can only appear when $u = c$. Also, Lemma (0.2) shows that if (u, ρ) is a traveling wave solution to (0.3), then $(u - c)^2 \in C^1(\mathbb{R})$. Now, consider the peaked function (see [2])

$$u = \chi + \sqrt{\chi^2 - c^2}, \quad \rho = \sqrt{-cK_1} \left(1 + \sqrt{\frac{\chi + c}{\chi - c}} \right), \quad \chi = -(c + K_1) \cosh(x - ct) + K_1,$$

where $K_1 = -\frac{1}{4}\kappa$, $K_1 < 0$ and $c > |K_1| > 0$. Away from its non-smooth point, u is a solution to (0.3). However, it is clear that u is not smooth at $\xi = 0$ and $u(0) = -c$. Furthermore, $(u - c)^2 \notin C^1(\mathbb{R})$ because

$$\lim_{\xi \downarrow 0} ((u - c)^2)' - \lim_{\xi \uparrow 0} ((u - c)^2)' = -8c\sqrt{c(c + K_1)},$$

where $\xi = x - ct$. Therefore, u is not a distributional solution to (0.3) even though it superficially looks like a peakon solution (see figure 3).




Figure 3: (u on the left and ρ on the right) This pair is not a distributional traveling wave solution of (0.3). $c = 2$ and $K_1 = -1$.

Acknowledgment. I would like to thank professor J. Szmigielski for suggesting the problem and tremendously helpful comments.

References

- [1] Roberto Camassa and Darryl D. Holm, *An integrable shallow water equation with peaked solitons*, arXiv:patt-sol/9305002 v1 13 May 1993
- [2] M. Cheng, S. Liu, Y. Zhang *A Two-component Generalization of the Camassa-Holm Equation and Its Solutions*, Letters in Mathematical Physics (2006) 75:1-15
- [3] I. M. Gel'fand and G. E. Shilov, *Generalized Functions*, Academic Press, New York and London, 1964

- [4] J. Lenells, *Traveling Wave Solutions of the Camassa-Holm Equation*, J. Differential Equations, 217 (2005) 393-430